arXiv:1202.2084v1 [quant-ph] 9 Feb 2012

Generation of GHZ entangled states of photons in multiple cavities via a superconucting qubit or an atom through resonant interaction

Chui-Ping Yang[1,2] and Siyuan Han[3]

[1]*Department of Physics, Hangzhou Normal University, Hangzhou, Zhejiang 310036, China*

[2]*State Key Laboratory of Precision Spectroscopy, Department of Physics,*
East China Normal University, Shanghai 200062, China and

[3]*Department of Physics and Astronomy, University of Kansas, Lawrence, Kansas 66045, USA*

(Dated: February 10, 2012)

We propose a method to generate a GHZ entangled state of n photons in n microwave cavities (or resonators) via resonant interaction to a single superconducting qubit. By performing local operations on a qubit (e.g., a solid-state qubit, an atom, etc.) placed in each cavity, the created GHZ states of n photons can be transferred to qubits for storage. The proposed scheme greatly reduces effect of decoherence since only resonant qubit-cavity interaction and resonant qubit-pulse interaction are involved, and no measurement is required. In addition, we show that the method can be applied to create a GHZ state of photons in multiple cavities via an atom through resonant interaction with no measurement needed.

PACS numbers: 03.67.Lx, 42.50.Dv, 85.25.Cp

I. INTRODUCTION

Entanglement is one of the most fascinating features of quantum mechanics and plays an important role in quantum communication and quantum information processing (QIP). During the past decade, experimental preparation of entanglement with eight photons via *linear optical devices* [1], eight ions [2], three spins [3], two atoms in microwave cavity QED [4], two atoms plus one cavity mode [5], or two excitons in a single quantum dot [6] has been reported. On the other hand, by having qubits coupled to a resonator or through a capacitive coupling, entangling two superconducting qubits [7-11] or three superconducting qubits [12,13] has been experimentally demonstrated. In addition, a tripartite entanglement with a superconducting qubit coupled to two microscopic two-level systems has been reported recently [14].

Over the past ten years, there has been much interest in quantum information processing with superconducting qubits in circuit QED. A cavity or resonator acts as a quantum bus which can mediate long-distance, fast interaction between distant superconducting qubits [15-18]. Theoretically, it was predicted earlier that the strong coupling limit can readily be realized with superconducting flux qubits [19] or charge qubits [17]. Moreover, the strong coupling limit between the cavity field and superconducting qubits has been experimentally demonstrated [20,21]. Based on circuit QED, a large number of theoretical schemes for creating entangled states with superconducting qubits in a cavity have been proposed [15,19,22-29]. In addition, two-qubit or three-qubit entangled states have been experimentally demonstrated with superconducting qubits coupled to a cavity or resonator [8-12]. All of these theoretical and experimental works are mainly focused on entanglement of superconducting qubits coupled to *one* cavity or resonator, which has paved the way for fundamental tests of quantum entanglement and nonlocality and made superconducting qubit circuit QED very attractive for quantum information processing.

Recently, attention has been shifting to entanglement generation of qubits or photons in multicavities because of its importance to scalable QIP. Within circuit QED, several theoretical proposals for generation of entangled photon Fock states of *two* resonators have been presented [30,31]. Moreover, by using a superconducting phase qubit coupled to two resonators, recent experimental demonstration of an entangled NOON state of photons in *two* superconducting microwave resonators has been reported [32].

In this paper, we focus on the preparation of GHZ (Greenberger-Horne-Zeilinger) entangled states of photons in *multiple* cavities. The GHZ entangled states are of great interest in the foundations of quantum mechanics and measurement theory, and significant in quantum information processing [33], quantum communication (e.g., cryptography) [34-36], error correction protocols [37], and high-precision spectroscopy [38].

In the following, we will propose an efficient method to generate a GHZ entangled state of n photons distributed over n microwave cavities or resonators coupled by a superconducting qubit (a.k.a. coupler) through resonant interaction. By local operations on a qubit (e.g., a solid-state qubit, an atom etc.) placed in each cavity, the created GHZ states of photons can be transferred to qubits for a long time storage and then can be transferred back to the photons once they are needed to be sent through quantum channels for implementing quantum communication or quantum information processing in a network.



FIG. 1: (Color online) Qubit-cavity resonant interaction. The cavity mode is resonant with the $|1\rangle \leftrightarrow |2\rangle$ transition of the qubit. g is the coupling constant between the cavity mode with the $|1\rangle \leftrightarrow |2\rangle$ transition. In (a), the level spacing between the two lowest levels is smaller than that between the upper two levels, this three-level structure applies to superconducting charge qubits and flux qubits [39]. In (b), the three-level structure is contrary to that in (a), and applies to superconducting phase qubits [39] as well as natural atoms. The qubit can be chosen such that the transition between the two lowest levels is forbidden due to the optical selection rules, weak via increasing the potential barrier between the two levels $|0\rangle$ and $|1\rangle$, or highly detuned (decoupled) from the cavity mode of each resonator by prior adjustment of the level spacings of the qubit.

As shown below, this proposal does not require measurement on the states of the coupler qubit or measurement on the cavity-mode photons for each cavity, and only requires resonant qubit-cavity interaction and resonant qubit-pulse interaction for each step of the operations. Thus, this proposal is simple to implement in experiments.

We stress that this proposal is quite general, and can be used to create a GHZ state of photons in multiple cavities or resonators when a different physical system, such as an atom or a quantum dot. In this paper, we show how to apply the method to generate a GHZ state of photons in multiple cavities using an atom.

This paper is organized as follows. In Sec. II, we briefly review the basic theory of resonant qubit-cavity and qubit-pulse interactions. In Sec. III, we show how to generate a GHZ state of n photons in n cavities coupled by a superconducting qubit. In Sec. IV, we give a brief discussion on the experimental issues and the experimental feasibility. In Sec. V, we show how to apply the method to create a GHZ state of n photons in n cavities via an atom. A concluding summary is given in Sec. VI.

II. BASIC THEORY

Consider a qubit A with three levels $|0\rangle$, $|1\rangle$, and $|2\rangle$ as depicted in Fig. 1. Suppose that the transition between the two levels $|1\rangle$ and $|2\rangle$ is resonant with the cavity mode. In the interaction picture, the interaction Hamiltonian of the qubit A and the cavity mode is given by

$$H = \hbar g(a^{+}\sigma_{12}^{-} + \text{H.c.}), \tag{1}$$

where a^{+} and a are the photon creation and annihilation operators of the cavity mode, g is the coupling constant between the cavity mode and the $|1\rangle \leftrightarrow |2\rangle$ transition of the qubit A, and $\sigma_{12}^{-} = |1\rangle \langle 2|$.

Based on the Hamiltonian (1), it can be easily found that the initial states $|2\rangle |0\rangle_c$ and $|1\rangle |1\rangle_c$ of the qubit A and the cavity mode evolve as follows

$$\begin{aligned} |2\rangle |0\rangle_c &\rightarrow -i\sin(gt) |1\rangle |1\rangle_c + \cos(gt) |2\rangle |0\rangle_c , \\ |1\rangle |1\rangle_c &\rightarrow \cos(gt) |1\rangle |1\rangle_c - i\sin(gt) |2\rangle |0\rangle_c . \end{aligned} \tag{2}$$

However, the states $|0\rangle |0\rangle_c$ and $|1\rangle |0\rangle_c$ remain unchanged under the Hamiltonian (1).

The coupling strength g may vary when the qubit A couples with different cavities or resonators. Therefore, in operations below, g is replaced by g_k to denote the coupling strength between the qubit A and cavity k ($k = 1, 2, ..., n$).

Let us now consider the case that qubit A is driven by a classical pulse of frequency ω. Suppose that the pulse is resonant with the transition between the two levels $|i\rangle$ and $|j\rangle$ of the qubit A, i.e., $\omega = \omega_{ji}$. Here, ω_{ji} is the $|i\rangle \leftrightarrow |j\rangle$ transition frequency of the qubit, and $|i\rangle$ is the lower energy level. The interaction Hamiltonian in the interaction picture is given by

$$H_I = \hbar \left(\Omega_{ij} e^{i\phi} |i\rangle \langle j| + \text{H.c.} \right), \tag{3}$$



FIG. 2: (Color online) (a) Diagram of a superconducting qubit A (a circle at the center) and n cavities. Each red dot represents a one-dimensional coplanar waveguide resonator which is coupled to the coupler qubit A through a capacitor, as shown in (b). (b) The diagram on the left side is equivalent to the diagram on the right side.

where Ω_{ij} is the Rabi frequency of the pulse and ϕ are the initial phase of the pulse. Based on the Hamiltonian (3), one can easily obtain the following state transformation

$$\begin{aligned} |i\rangle &\rightarrow \cos\Omega_{ij}t\,|i\rangle - ie^{-i\phi}\sin\Omega_{ij}t\,|j\rangle\,, \\ |j\rangle &\rightarrow \cos\Omega_{ij}t\,|j\rangle - ie^{i\phi}\sin\Omega_{ij}t\,|i\rangle\,, \end{aligned} \tag{4}$$

which can be completed rapidly by increasing Ω_{ij} (i.e., by increasing the intensity of the pulse).

In above, we have reviewed the basic theory of the qubit A resonantly interacting with the cavity mode or a pulse. The results presented above will be employed for the entanglement preparation discussed in next section.

III. GENERATION OF A N-PHOTON GHZ STATE VIA A SUPERCONDUCTING QUBIT

Consider n cavities $(1, 2, ..., n)$ each coupled to a superconducting coupler qubit A, which has three levels as depicted in Fig. 1. The three-level structure in Fig. 1(a) applies to superconducting charge qubits and flux qubits, and the one in Fig. 1(b) applies to phase qubits (see Ref. [39] and other relevant references cited in [39]). In addition, the three-level structure in Fig. 1(b) is also available in atoms. The coupler qubit A can be chosen such that the transition between the two lowest levels is forbidden due to the optical selection rules [40], weak via increasing the potential barrier between the two levels $|0\rangle$ and $|1\rangle$ [41-43], or highly detuned (decoupled) from the cavity mode of each resonator by prior adjustment of the level spacings of the qubit. Note that for superconducting qubits, the level spacings can be readily adjusted by varying external control parameters (e.g., gate voltage and/or magnetic flux applied to the superconducting charge qubits and/or phase qubits and flux qubits, see e.g. [41-44]).

Initially, the qubit A is decoupled from the cavities $(1, 2, ..., n)$, which can be realized by prior adjustment of each cavity frequency; the qubit A is prepared in the state $\left(|0\rangle_A + |2\rangle_A\right)/\sqrt{2}$; and each cavity is in a vacuum state, i.e., $|0\rangle_{c,i}$ for cavity i $(i = 1, 2, ..., n)$. The operations for realizing a GHZ state of n photons in the n cavities are listed below:

Step *1*: Adjust the frequency ω_{c1} of cavity 1 such that it is resonant with the $|1\rangle \leftrightarrow |2\rangle$ transition of qubit A. After an interaction time $t_{1a} = \pi/(2g_1)$, the state $|0\rangle_A |0\rangle_{c1}$ remains unchanged while the state $|2\rangle_A |0\rangle_{c1}$ changes to

$-i\left|1\right\rangle_A\left|1\right\rangle_{c1}$, as described by Eq. (2). Then, adjust the frequency of cavity 1 such that it is decoupled from qubit A and then apply a pulse (with frequency $\omega=\omega_{21}$, initial phase $\phi=\pi$, and duration $t_{1b}=\pi/\left(2\Omega_{21}\right)$) to qubit A, to transform the state $\left|1\right\rangle_A$ to $i\left|2\right\rangle_A$, as described by Eq. (4).

It can be seen that after this step of operation, the initial state $\left(\left|0\right\rangle_A+\left|1\right\rangle_A\right)\prod_{i=1}^{n}\left|0\right\rangle_{ci}$ of the whole system changes to (here and below a normalization factor is omitted for simplicity)

$$\left(\left|0\right\rangle_A\left|0\right\rangle_{c1}+\left|2\right\rangle_A\left|1\right\rangle_{c1}\right)\prod_{i=2}^{n}\left|0\right\rangle_{ci}. \tag{5}$$

Step *2*: Adjust the frequency ω_{c2} of cavity 2 to have it resonant with the $\left|1\right\rangle\leftrightarrow\left|2\right\rangle$ transition of qubit A for an interaction time $t_{2a}=\pi/(2g_2)$. As a result, the state $\left|2\right\rangle_A\left|0\right\rangle_{c2}$ changes to $-i\left|1\right\rangle_A\left|1\right\rangle_{c2}$ while the state $\left|0\right\rangle_A\left|0\right\rangle_{c2}$ remains unchanged. Then, adjust the frequency of cavity 2 to have it decoupled from qubit A and then apply a pulse (with frequency $\omega=\omega_{21}$, initial phase $\phi=\pi$, and duration $t_{2b}=\pi/\left(2\Omega_{21}\right)$) to qubit A, to transform the state $\left|1\right\rangle_A$ to $i\left|2\right\rangle_A$.

It can be seen that after this step of operation, the state (5) changes to

$$\left(\left|0\right\rangle_A\left|0\right\rangle_{c1}\left|0\right\rangle_{c2}+\left|2\right\rangle_A\left|1\right\rangle_{c1}\left|1\right\rangle_{c2}\right)\prod_{i=3}^{n}\left|0\right\rangle_{ci}. \tag{6}$$

Step *3* to Step (*n-2*): The procedure for each step of operations from step *3* to step (*n-2*) is the same as step *1* or step *2*. Namely, for step k ($k=3,4,...,n-2$), we adjust the frequency ω_{ck} of cavity k to have it resonant with the $\left|1\right\rangle\leftrightarrow\left|2\right\rangle$ transition of qubit A. After an interaction time $t_{ka}=\pi/(2g_k)$, the state $\left|2\right\rangle_A\left|0\right\rangle_{ck}$ changes to $-i\left|1\right\rangle_A\left|1\right\rangle_{ck}$ while the state $\left|0\right\rangle_A\left|0\right\rangle_{ck}$ remains unchanged. Then, adjust the frequency of cavity k to have it decoupled from qubit A and then apply a pulse (with frequency $\omega=\omega_{21}$, initial phase $\phi=\pi$, and duration $t_{kb}=\pi/\left(2\Omega_{21}\right)$) to qubit A, to transform the state $\left|1\right\rangle_A$ to $i\left|2\right\rangle_A$.

One can verify that after the operations of step *3* to step (*n-2*), the state (6) changes to

$$\left(\left|0\right\rangle_A\prod_{i=1}^{n-2}\left|0\right\rangle_{ci}+\left|2\right\rangle_A\prod_{i=1}^{n-2}\left|1\right\rangle_{ci}\right)\left|0\right\rangle_{c(n-1)}\left|0\right\rangle_{cn}. \tag{7}$$

Step (*n-1*): Adjust the frequency $\omega_{c(n-1)}$ of cavity $(n-1)$ to have it resonant with the $\left|1\right\rangle\leftrightarrow\left|2\right\rangle$ transition of qubit A for an interaction time $t_{(n-1)a}=\pi/(2g_{n-1})$. As a result, the state $\left|2\right\rangle_A\left|0\right\rangle_{c(n-1)}$ changes to $-i\left|1\right\rangle_A\left|1\right\rangle_{c(n-1)}$ while the state $\left|0\right\rangle_A\left|0\right\rangle_{c(n-1)}$ remains unchanged. Then, adjust the frequency of cavity $n-1$ to have it decoupled from qubit A. Next, apply a pulse (with frequency $\omega=\omega_{20}$, initial phase $\phi=-\pi/2$, and duration $t_{(n-1)b}=\pi/\left(2\Omega_{20}\right)$) to qubit A, to transform the state $\left|0\right\rangle_A$ to $\left|2\right\rangle_A$; and then apply a pulse (with frequency $\omega=\omega_{21}$, initial phase $\phi=\pi/2$, and duration $t_{(n-1)c}=\pi/\left(2\Omega_{21}\right)$) to qubit A, to transform the state $\left|1\right\rangle_A$ to $-\left|2\right\rangle_A$ while the state $\left|2\right\rangle_A$ to $\left|1\right\rangle_A$.

It can be found that after this step of operation, the following transformation is obtained

$$\begin{array}{ccc}\left|0\right\rangle_A\left|0\right\rangle_{c(n-1)} & \overset{\text{Step n-1}}{\rightarrow} & \left|1\right\rangle_A\left|0\right\rangle_{c(n-1)} \\ \left|2\right\rangle_A\left|0\right\rangle_{c(n-1)} & & i\left|2\right\rangle_A\left|1\right\rangle_{c(n-1)}\end{array}. \tag{8}$$

Based on Eq. (8), one can see that the state (7) changes to

$$\left(\left|1\right\rangle_A\prod_{i=1}^{n-1}\left|0\right\rangle_{ci}+i\left|2\right\rangle_A\prod_{i=1}^{n-1}\left|1\right\rangle_{ci}\right)\left|0\right\rangle_{n}. \tag{9}$$

Step n: Adjust the frequency ω_{cn} of cavity n such that it is resonant with the $\left|1\right\rangle\leftrightarrow\left|2\right\rangle$ transition of qubit A. After an interaction time $t_n=\pi/(2g_n)$, the state $\left|2\right\rangle_A\left|0\right\rangle_{cn}$ changes to $-i\left|1\right\rangle_A\left|1\right\rangle_{cn}$ while the state $\left|1\right\rangle_A\left|0\right\rangle_{cn}$ remains unchanged. Then, adjust the frequency of cavity n to have it decoupled from qubit A.

It can be seen that after this step of operation, the state (9) becomes

$$\left|1\right\rangle_A\left(\prod_{i=1}^{n}\left|0\right\rangle_{ci}+\prod_{i=1}^{n}\left|1\right\rangle_{ci}\right). \tag{10}$$

The result (10) shows that the n cavities are prepared in a n-photon GHZ state $\prod_{i=1}^{n}\left|0\right\rangle_{ci}+\prod_{i=1}^{n}\left|1\right\rangle_{ci}$, while the qubit A is disentangled from all cavities, after the above operations.

In principle, the method presented above can be used to create a GHZ state of n photons in n cavities. However, it should be pointed out that in practice since g_i is inversely proportional to n, the number of cavities coupled to qubit A is limited to maintain sufficiently strong couplings.

As discussed above, the present method requires rapid adjustments of the cavity frequency during entanglement preparation. Note that for superconducting microwave cavities, the cavity frequency can be fast tuned (e.g., in less than a few nanoseconds for a superconducting transmission line resonator [45]).

Finally, it should be mentioned that in principle, the method can also be realized with cavities of different resonant frequencies by rapid tuning of level spacing ω_{21} of the coupler qubit.

IV. DISCUSSION

Let us now give some discussions on the relevant experimental issues for the entanglement preparation. For the method to work:

(i) The total operation time τ, given by

$$\tau=\sum_{j=1}^{n}\pi/(2g_j)+(n-1)\,\pi/\left(2\Omega_{21}\right)+\pi/\left(2\Omega_{20}\right)+2nt_d \tag{11}$$

(where t_d is the typical time required for adjusting the cavity mode frequency), needs to be much shorter than the energy relaxation time T_1 and dephasing time T_2 of the level $|2\rangle$ of qubit A, such that decoherence caused due to energy relaxation time and dephasing time of qubit A is negligible during the entire operation. Note that the energy relaxation time T_1' and dephasing time T_2' of the level $|1\rangle$ of qubit A are longer than T_1 and T_2, respectively.

(ii) For cavities $(1,2,...,n)$, the lifetime of the cavity mode is given by $T_{cav}^{i}=\left(Q_i/2\pi\nu_{c,i}\right)/\overline{n}_i$, where Q_i and $\overline{n}_i$ are the (loaded) quality factor and the average photon number of the cavity i, respectively. For n cavities, the lifetime of the cavity modes is given by

$$T_{cav}=\frac{1}{n}\min\{T_{cav}^{1},T_{cav}^{2},...,T_{cav}^{n}\}, \tag{12}$$

which should be much longer than τ, such that the cavity decay is negligible during the entire operation.

These requirements can in principle be realized, since one can: (i) reduce the qubit-cavity interaction time by increasing the qubit-cavity coupling constant g_j, (ii) shorten t_d by a rapid adjustment of the cavity mode frequency, (iii) increase T_{cav} by employing high-Q cavities, and (iv) utilize qubit A with long T_1 and T_2 times.

For the sake of definitiveness, let us consider the experimental feasibility of generating a GHZ state of 10 photons in the ten cavities. Without loss of generality, we consider a case that the coupling contant of qubit A with each cavity is identical, i.e, $g_1=g_2=...=g_{10}=g$. We assume $g/2\pi\sim 150$ MHz, which could be reached for a superconducting qubit coupled to a one-dimensional standing-wave CPW (coplanar waveguide) transmission line resonator [12]. With the choice of $\Omega_{21}\sim\Omega_{20}\sim 2g$ and $t_d\sim 3$ ns [45], a simple calculation gives $\tau\sim 85$ ns, which is much shorter than $\min\{T_1,T_2\}\sim 5$ μs for $T_1\sim T_1'/2$ and $T_2\sim T_2'/2$. Note that T_1' and T_2' can be made to be on the order of ~ 10 μs for the state of art superconducting qubits at this time [46].

Superconducting CPW transmission line resonators with a loaded quality factor $Q\sim 10^6$ have been experimentally demonstrated [47,48]. Without loss of generality, consider ten resonators each with a center frequency ~ 5 GHz [7] and a quality factor $\sim 5\times 10^5$. From the discussion given above, it can be seen that each cavity was occupied by a single photon during the GHZ-state preparation. Thus, we have $\min\{T_{cav}^{1},T_{cav}^{2},...,T_{cav}^{10}\}\sim 16$ μs, resulting in $T_{cav}\sim 1.6$ μs for $n=10$ for the present case, which is much longer than τ. The analysis given here shows that generating a GHZ state of 10 photons in ten cavities might be possible within the present circuit QED technique.

V. GENERATION OF A N-PHOTON GHZ STATE USING AN ATOM

During the past decade, much attention has been paid to the generation of highly entangled states with atomic systems. Two-atom entangled states and three-particle GHZ entangled states (with two atoms plus one cavity mode) have been experimentally demonstrated in microwave cavity QED [4,5]. In addition, based on cavity QED, numerous theoretical proposals have been presented for entangling atoms coupling to the mode (s) of a single cavity [49] and atoms in two or more cavities [50]. In principle, an entangled state of n photons in n cavities ($n\geq 2$) can be created, by first preparing an n-atom entangled state using the previous proposals [49,50], and then transferring the prepared n-atom entangled states onto n photons in the n cavities via the state transfer from an atom to a photon in a cavity. In the following, we will present an alternative way to implement an n-photon GHZ state, which, as shown below, does not require prior preparation of atomic entangled states. The scheme presented here is actually a generalization of the method described in Sec. II to GHZ-state generation of photons in multiple cavities through an atom.



FIG. 3: (Color online) Diagram of n cavities and an atom A (a red dot). Before atom A arrives in cavity $n-1$, the atom A is moved into each cavity to have it interacting with the cavity mode for a given time and then moved out of the cavity to have it addressed by a classical pulse with frequency equal to the transition frequency ω_{21} of the atom A (see the pink-color frame with an arrow). When the atom A leaves the cavity $n-1$, two pulses are applied to the atom A. The first pulse has a frequency equal to the transition frequency ω_{20} (see the blue-color frame with an arrow) while the second pulse has a frequency equal to the transition frequency ω_{21} of the atom A.

Consider n cavities $(1,2,...,n)$ and an atom A with three levels as depicted in Fig. 1. The atom A is initially prepared in the state $\left(|0\rangle_A+|2\rangle_A\right)/\sqrt{2}$ and each cavity is in a vacuum state, i.e., $|0\rangle_{c,i}$ for cavity i $(i=1,2,...,n)$. In addition, assume that the cavity mode of each cavity is resonant with the $|1\rangle\leftrightarrow|2\rangle$ transition but highly detuned (decoupled) from the transition between any other two levels of the atom A. The procedure for generating a GHZ state of n photons in the n cavities is illustrated in Fig. 2. The operations shown in Fig. 2 are as follows:

Step *1*: Move atom A into cavity 1 for an interaction time $t_{1a}=\pi/(2g_1)$. As a result, the state $|0\rangle_A|0\rangle_{c1}$ remains unchanged while the state $|2\rangle_A|0\rangle_{c1}$ changes to $-i|1\rangle_A|1\rangle_{c1}$, as described as Eq. (2). Then move atom A out of the cavity 1 and apply a classical pulse (with frequency $\omega=\omega_{21}$, initial phase $\phi=\pi$, and duration $t_{1b}=\pi/(2\Omega_{21})$) to atom A, to transform the state $|1\rangle_A$ to $i|2\rangle_A$, as described by Eq. (4). It can be seen that after this step of operation, the initial state $\left(|0\rangle_A+|1\rangle_A\right)\prod_{i=1}^{n}|0\rangle_{ci}$ of the whole system changes to the state given in Eq. (5).

Step *2*: Move atom A into cavity 2 for an interaction time $t_{2a}=\pi/(2g_2)$. As a result, the state $|2\rangle_A|0\rangle_{c2}$ changes to $-i|1\rangle_A|1\rangle_{c2}$ while the state $|0\rangle_A|0\rangle_{c2}$ remains unchanged. Then move atom A out of the cavity 2 and apply a pulse (with frequency $\omega=\omega_{21}$, initial phase $\phi=\pi$, and duration $t_{2b}=\pi/(2\Omega_{21})$) to atom A, to transform the state $|1\rangle_A$ to $i|2\rangle_A$. It can be seen that after this step of operation, the state (5) changes to the state (6).

Step *3* to Step (*n-2*): The procedure for each step of operations from step *3* to step (*n-2*) is the same as either step *1* or step *2*. Namely, for step k $(k=3,4,...,n-2)$, we move atom A into cavity k for an interaction time $t_{ka}=\pi/(2g_k)$, such that the state $|2\rangle_A|0\rangle_{ck}$ changes to $-i|1\rangle_A|1\rangle_{ck}$ while the state $|0\rangle_A|0\rangle_{ck}$ remains unchanged. Then, move atom A out of cavity k and apply a pulse (with frequency $\omega=\omega_{21}$, initial phase $\phi=\pi$, and duration $t_{kb}=\pi/(2\Omega_{21})$) to atom A, to transform the state $|1\rangle_A$ to $i|2\rangle_A$. It is easy to verify that after the operations of step *3* to step (*n-2*), the state (6) changes to the state (7).

Step *(n-1)*: Move atom A into cavity $(n-1)$ for an interaction time $t_{(n-1)a}=\pi/(2g_{n-1})$, such that the state $|2\rangle_A|0\rangle_{c(n-1)}$ changes to $-i|1\rangle_A|1\rangle_{c(n-1)}$ while the state $|0\rangle_A|0\rangle_{c(n-1)}$ remains unchanged. Then move atom A out of the cavity $n-1$ and apply a pulse (with frequency $\omega=\omega_{20}$, initial phase $\phi=-\pi/2$, and duration $t_{(n-1)b}=\pi/(2\Omega_{20})$) to atom A, to transform the state $|0\rangle_A$ to $|2\rangle_A$; and then apply a pulse (with frequency $\omega=\omega_{21}$, initial phase $\phi=\pi/2$, and duration $t_{(n-1)c}=\pi/(2\Omega_{21})$) to atom A, to transform the state $|1\rangle_A$ to $-|2\rangle_A$ while the state $|2\rangle_A$ to $|1\rangle_A$. One can check that after this step of operation, the following transformation given in Eq. (8) is achieved. Based on Eq. (8), one can see that the state (7) changes to the state (9).

Step n: Move atom A into cavity n for an interaction time $t_n=\pi/(2g_n)$, such that the state $|2\rangle_A|0\rangle_{cn}$ changes to $-i|1\rangle_A|1\rangle_{cn}$ while the state $|1\rangle_A|0\rangle_{cn}$ remains unchanged. After atom A interacting with cavity n for a time t_n, move atom A out of the cavity n such that it is decoupled from the cavity n. It can be seen that after this step of operation, the state (9) becomes the state (10), which shows that a GHZ state $\prod_{i=1}^{n}|0\rangle_{ci}+\prod_{i=1}^{n}|1\rangle_{ci}$ of n photons in n cavities is created while the atom A is disentangled from each of the cavities, after the above process.

It should be mentioned that the atom A can be trapped in a linear trap [51], inside an optical lattice [52], or on top of an atomic chip [53]; and then it can be moved into or out of a cavity, by moving the lattice or by applying an electric field [54]. The approach for trapping and moving atoms has been employed in the earlier work for quantum computing with atoms in cavity QED [54-57]. Alternatively, the atom A can be sent through a cavity to have it interacting with the cavity for a given interaction time, by choosing the atomic velocity appropriately [58].

From the above description, it is concluded that the present scheme has the following advantages:

(i) Only one atom is needed for the entanglement preparation of photons in multiple cavities;

(i) Neither measurement on the states of the atom A nor measurement on the cavity photons is needed;

(ii) No adjustment of the atomic level spacings is needed during the entire operation; and also

(iii) No adjustement of the cavity mode frequency is required during the entire operation.

For our scheme to work, the total operation time τ given in Eq. (12) (where τ_d is now a typic time for moving atom A into or out of a cavity) should be much smaller than the lifetime of the cavity modes of n cavities, i.e., T_{cav} given in Eq. (13), so that the cavity dissipation is negligible during the entire operation. In addition, the τ needs to be much shorter than the spontaneous emission time T_r of the level $|2\rangle$ of atom A, such that decoherence caused due to the spontaneous emission of atom A is negligible. Note that the spontaneous emission time of the level $|1\rangle$ of atom A is longer than T_r.

To investigate the experimental feasibility of this scheme, let us consider preparation of a GHZ state for 10 photons in ten cavities using a single Rydberg atom. Without loss of generaility, we assume $g_1 = g_2 = ... = g_{10} = g$. The atom A is chosen as a Rydberg atom with principal quantum numbers 50 and 51 (respectively corresponding to the levels $|1\rangle$ and $|2\rangle$). For the Rydberg atom chosen here, the $|1\rangle \leftrightarrow |2\rangle$ transition frequency is $\omega_0/2\pi \sim 51.1$ GHz, the energy relaxation time of the level $|2\rangle$ is $T_r \sim 3 \times 10^{-2}$ s, and the coupling constant is $g = 2\pi \times 50$ kHz [59]. With the choice of $t_d \sim 1\mu$s and $\Omega_{21} \sim \Omega_{20} \sim 10g$, we have $\tau \sim 7.5 \times 10^{-5}$ s, much shorter than T_r.

In the present case, the cavity mode frequency of each cavity is ~ 51.1 GHz. One can see from the above discussion that each cavity was occupied by a single photon during the GHZ-state preparation. For a cavity with $Q = 10^{10}$, we have $\min\{T^1_{cav}, T^2_{cav}, ..., T^{10}_{cav}\} \sim 3.1 \times 10^{-2}$ s, resulting in $T_{cav} \sim 3.1 \times 10^{-3}$ s for $n = 10$. which is much longer than τ. Note that cavities with a high $Q \sim 10^{12}$ have been reported [60]. Thus, generating a GHZ state of 10 photons in ten cavities with assistance of an atom is possible within the present cavity QED technique.

By Using *linear optics elements* and *single photon detectors*, many schemes for creating photon entangled states have also been proposed [61]; and experimental realization of an eight-photon GHZ state [1] and a three-photon W state [62] has been reported. However, this type of approaches is much more difficult to implement than cavity QED for hybrid systems consisting of photons and matter qubits of nature made and/or engineered. The present work represents a significant advancement in circuit and atom QED because it provides a simple and fast approach for deterministically creating a multi-photon GHZ state, which does not require measurement or detection on photons.

We noticed that two previous works [63,64] are relevant to ours. Ref. [63] presents a scheme for preparation of a GHZ-type entangled *coherent state* of n cavities by having an atom interacts with each of the cavities dispersively and then measuring the state of the atom. We realize that a GHZ entangled Fock state of photons in multiple cavities can in principle be generated using the same procedure described in [63]. However, the method has the following drawbacks: (i) the operation is rather slow because of the dispersive atom-cavity interaction, (ii) a measurement on the state of the atom is required, and (iii) since the prepared GHZ state depends on the measurement outcome on the atomic states, the GHZ-state preparation is not deterministic. In contrast, our proposal mitigates these problems effectively: the operation is much faster because of the resonant atom-cavity interactions; there is no need to measure the state of the atom; and the generation of the GHZ state is deterministic. Ref. [64] proposes a method for preparing a cluster state of photons in n cavities via resonant atom-cavity interactions. However, our proposal is significantly different from that of [64]. First, we focus on preparing a GHZ entangled Fock state of photons in multiple cavities. Second, an n-qubit cluster state cannot be transformed into a GHZ state (for $n > 3$) [65]. Last, the method proposed in [64] requires an atom to interact with *two* classical pulses after it leaves each cavity (except the final one) while our proposal only requires the atom interacting with *one* classical pulse after it exits each cavity (except the final one).

After a thorough search, we found that three schemes [66-68] were previously proposed for implementing the GHZ state of photons in n cavities by sending an atom through n cavities. However, these schemes require measuring the state of the atom and/or using n levels of the atom (i.e., the number of the atomic levels used needs to be equal to the number of the cavities).

VI. CONCLUSION

We have presented a method to generate a GHZ state of n photons in n cavities coupled by a superconducting qubit. By local operations on a qubit (e.g., an atom or a solid-state qubit) placed in each cavity, the created GHZ states of photons can be transferred to qubits for the storage for a long time. This proposal is easy to be implemented in experiments since only resonant qubit-cavity interaction and resonant qubit-pulse interaction are needed, and no measurement is required. In addition, we have shown how to apply the present method to create a GHZ state of n photons in n cavities via an atom. We note that neither adjusting the atomic level spacings nor adjusting the cavity mode frequency is needed during the entire operation and only one atom is needed for the entanglement preparation of photons in multiple cavities. In addition, our analysis shows that generating a GHZ state of ten photons in ten cavities by a coupler superconducting qubit or an atom is possible within the present experimental technique. Finally, it should be mentioned that this proposal is quite general, which can be applied to create a GHZ state of photons in multiple cavities or resonators, when the couper qubit is a different physical system, such as a quantum dot or an NV center.

ACKNOWLEDGMENTS

C.P. Yang was supported in part by the National Natural Science Foundation of China under Grant No. 11074062, the Zhejiang Natural Science Foundation under Grant No. Y6100098, the Open Fund from the SKLPS of ECNU, and the funds from Hangzhou Normal University. S. Han was supported in part by DMEA.

---

[1] Y. F. Huang *et al.*, Nature Communications **2**, 546 (2011); X. C. Yao *et al.*, arXiv:1105.6318.
[2] H. Häffner *et al.*, Nature (London) **438**, 643 (2005).
[3] D. M. Greenberger, M. A. Horne, and A. Zeilinger, Going beyond Bells theorem. In Kafatos, M. (ed.) *Bells theorem, quantum theory and conceptions of the universe* (Kluwer Academic, Dordrecht, 1989).
[4] E. Hagley *et al.*, Phys. Rev. Lett. 79, 1 (1997); S. Osnaghi *et al.*, *ibid.* **87**, 037902 (2001).
[5] A. Rauschenbeutel *et al.*, Science **288**, 2024 (2000).
[6] G. Chen, N. H. Bonadeo, D. G. Steel, D. Gammon, D. S. Katzer, D. Park, and L. J. Sham, Science **289**, 1906 (2000).
[7] M. Steffen *et al.*, Science **313**, 1423 (2006).
[8] P. J. Leek *et al.* Using sideband transitions for two-qubit operations in superconducting circuits. Phys. Rev. B 79, 180511(R) (2009).
[9] L. DiCarlo *et al.*, Nature **460**, 240 (2009).
[10] M. Ansmann *et al.*, Nature **461**, 504506 (2009)
[11] J. M. Chow *et al.*, arXiv:0908.1955.
[12] L. DiCarlo, M. D. Reed, L. Sun, B. R. Johnson, J. M. Chow, J. M. Gambetta, L. Frunzio, S. M. Girvin, M. H. Devoret, and R. J. Schoelkopf, Nature (London) **467**, 574 (2010).
[13] M. Neeley *et al.*, Nature (London) **467**, 570 (2010).
[14] G. Sun, X. Wen, B. Mao, J. Chen, Y. Yu, P. Wu, and S. Han, Nature Communications **1**, 51 (2010).
[15] C. P. Yang, Shih-I Chu, and S. Han, Phys. Rev. A **67**, 042311 (2003).
[16] J. Q. You and F. Nori, Phys. Rev. B **68**, 064509 (2003).
[17] A. Blais *et al.*, Phys. Rev. A **69**, 062320 (2004).
[18] J. Majer *et al.*, Nature (London) **449**, 443 (2007).
[19] C. P. Yang, Shih-I Chu, and S. Han, Phys. Rev. Lett. **92**, 117902 (2004).
[20] A. Wallraff *et al.*, Nature (London) **431**, 162 (2004).
[21] I. Chiorescu *et al.*, Nature (London) **431**, 159 (2004).
[22] C. P. Yang and S. Han, Phys. Rev. A **70**, 062323 (2004).
[23] C. P. Yang, Shih-I Chu, and S. Han, J. Phys.: Condens. Matter **16**, 1907 (2004).
[24] S. L. Zhu, Z. D. Wang, and P. Zanardi, Phys. Rev. Lett **94**, 100502 (2005).
[25] K. H. Song, Z. W. Zhou, and G. C. Guo, Phys. Rev. A **71**, 052310 (2005)
[26] X. L. Zhang, K. L. Gao, and M. Feng, Phys. Rev. A **74**, 024303 (2006);
[27] Z. J. Deng, K. L. Gao, and M. Feng, Phys. Rev. A **74**, 064303 (2006).
[28] K. H. Song, S. H. Xiang, Q. Liu, and D. H. Lu, Phys. Rev. A **75**, 032347 (2007).
[29] C. P. Yang, Phys. Rev. A **83**, 062302 (2011).
[30] M. Mariantoni *et al.*, Phys. Rev. B **78**, 104508 (2008).
[31] F. W. Strauch *et al.*, Phys. Rev. Lett. **105**, 050501 (2010).
[32] H. Wang *et al.*, Phys. Rev. Lett. **106**, 060401 (2011).
[33] M. Hillery, V. Buzek, and A. Berthiaume, Phys. Rev. A **59**, 1829 (1999).
[34] S. Bose, V. Vedral, and P. L. Knight, Phys. Rev. A **57**, 822 (1998).
[35] R. Cleve, D. Gottesman, and H. K. Lo, Phys. Rev. Lett. **83**, 648 (1999); V. Scarani and N. Gisin, *ibid.* **87**, 117901 (2001); G. A. Durkin, C. Simon, and D. Bouwmeester, *ibid.* **88**, 187902 (2002).
[36] C. P. Yang, Shih-I Chu, and S. Han, Phys. Rev. A **70**, 022329 (2004)
[37] D. P. DiVincenzo and P. W. Shor, Phys. Rev. Lett. **77**, 3260 (1996); J. Preskill, Proc. R. Soc. London, Ser. A **454**, 385 (1998).
[38] J. J. Bollinger, W. M. Itano, D. J. Wineland, and D. J. Heinzen, Phys. Rev. A **54**, 4649 (1996); S. F. Huelga, C. Macchiavello, T. Pellizzari, A. K. Ekert, M. B. Plenio, and J. I. Cirac, Phys. Rev. Lett. **79**, 3865 (1997).
[39] C. P. Yang, S. B. Zheng, and F. Nori, Phys. Rev. A **82**, 062326 (2010).
[40] Y. X. Liu, J. Q. You, L. F.Wei, C. P. Sun, and F. Nori, Phys. Rev. Lett. 95, 087001 (2005).
[41] S. Han, J. Lapointe, and J. E. Lukens, Single-Electron Tunneling and Mesoscopic Devices (Springer-Verlag, Berlin-Heidelberg, 1991), Vol. 31, pp. 219222.
[42] J. Clarke and F. K.Wilhelm, Nature (London) 453, 1031 (2008).
[43] M. Neeley *et al.*, Nat. Phys. 4, 523 (2008).
[44] J. Q. You and F. Nori, Phys. Today 58, 42 (2005).
[45] M. Sandberg, C.M.Wilson, F. Persson, T. Bauch, G. Johansson, V. Shumeiko, T. Duty, and P. Delsing, Appl. Phys. Lett. 92, 203501 (2008).
[46] J. Bylander *et al.*, Nature Phys. **7**, 565 (2011).
[47] W. Chen, D. A. Bennett, V. Patel, and J. E. Lukens, Supercond. Sci. Technol. **21**, 075013 (2008).

[48] P. J. Leek *et al.*, Phys. Rev. Lett. 104, 100504 (2010).

[49] J. I. Cirac and P. Zoller, Phys. Rev. A 50, R2799 (1994); C. C. Gerry, *ibid.* **53**, 2857 (1996); M. B. Plenio, S. F. Huelga, A. Beige, and P. L. Knight, *ibid.* **59**, 2468 (1999); A. S. Søensen and Klaus Mømer, *ibid.* **66**, 022314 (2002); G. P. Guo, C. F. Li, J. Li, and G. C. Guo, *ibid.* **65**, 042102 (2002); Y. F. Xiao, X. B. Zou, and G. C. Guo, *ibid.* **75**, 012310 (2007); L. M. Duan and H. J. Kimble, *ibid.* **90**, 253601 (2003); S. B. Zheng, Phys. Rev. Lett. 87, 230404 (2001); Phys. Rev. A 66, 060303 (2002).

[50] J. I. Cirac, P. Zoller, H. J. Kimble, and H. Mabuchi, Phys. Rev. Lett. 78, 3221 (1997); S. J. van Enk, J. I. Cirac, and P. Zoller, *ibid.* **78**, 4293 (1997); L. M. Duan and H. J. Kimble, *ibid.* **90**, 253601 (2003); X. L. Feng, Z. M. Zhang, X. D. Li, S. Q. Gong, and Z. Z. Xu, *ibid.* **90**, 217902 (2003); P. Xue, C. Han, B. Yu, X. M. Lin, and G. C. Guo, Phys. Rev. A **69**, 052318 (2004); J. H. An, M. Feng, and C. H. Oh, *ibid.* **032303** (2009); X. L. Zhang, K. L. Gao, and M. Feng, *ibid.* 75, 034308 (2007); P. Peng and F. L. Li, *ibid.* **75**, 062320 (2007); J. Lee, J. Park, S. M. Lee, H. W. Lee, and A. H. Khosa, *ibid.* **77**, 032327 (2008); J. Cho, D. G. Angelakis, S. Bose , *ibid.* **78**, 022323 (2008); S. B. Zheng, C. P. Yang, and F. Nori, *ibid.* **82**, 042327 (2010).

[51] G.R. Guthöhrlein, M. Keller, K. Hayasaka, W. Lange, H. Walther, Nature **414**, 49 (2001).

[52] S. Kuhr, W. Alt, D. Schrader, M. Müller, V. Gomer, D. Meschede, Science **293**, 278 (2001); J. Cho, Phys. Rev. Lett. **99**, 020502 (2007); H. Zoubi, H. Ritsch, Phys. Rev. A **76**, 013817 (2007); D. Jaksch, quant-ph/0407048.

[53] P. Horak, B.G. Klappauf, A. Haase, R. Folman, J. Schmiedmayer, P. Domokos, E.A. Hinds, Phys. Rev. A **67**, 043806 (2003).

[54] A. Beige, D. Braun, B. Tregenna, P.L. Knight, Phys. Rev. Lett. **85**, 1762 (2000).

[55] B. Tregenna, A. Beige, P. L. Knight, Phys. Rev. A **65** 032305 (2002).

[56] A. Beige, H. Cable, P. L. Knight, Proc. SPIE **5111** 370 (2003).

[57] L. M. Duan, B. Wang, and H. J. Kimble, Phys. Rev. A **72**, 032333 (2005).

[58] J. I. Cirac and P. Zoller, Phys. Rev. A 50, R2799 (1994); C. C. Gerry, *ibid.* **53**, 2857 (1996); S. B. Zheng, Phys. Rev. Lett. 87, 230404 (2001).

[59] M. Brune *et al.*, Phys. Rev. Lett. **77** 4887 (1996); S. Osnaghi *et al.*, *ibid.* **87**, 037902 (2001).

[60] G. Rempe, F. SchmidtKaler, and H. Walther, Phys. Rev. Lett. **64**, 2783 (1990).

[61] H. Lee, P. Kok, N. J. Cerf, and J. P. Dowling, Phys. Rev. A **65**, 030101(R) (2002); J. Fiurásek **65**, 053818 (2002); X. B. Zou, K. Pahlke, and W. Mathis, *ibid.* **66**, 014102 (2002); *ibid.* **66**, 044302 (2002); T. Yamamoto, K. Tamaki, M. Koashi, and N. Imoto, *ibid.* **66**, 064301 (2002); H. Jeong and N. Ba An, *ibid.* **74**, 022104 (2006); Y. Sagi, *ibid.* **68**, 042320 (2003).

[62] M. Eibl *et al.*, Phys. Rev. Lett. **92**, 077901 (2004).

[63] C. C. Gerry, Phys. Rev. A *54*, R2529 (1996).

[64] X. B. Zou and W. Mathis, Phys. Rev. A **72**, 013809 (2005).

[65] H. J. Briegel and R. Raussendorf, Phys. Rev. Lett. **86**, 910 (2001).

[66] Z. M. Zhan, J. H. Li, W. X. Yang, and W. B. Li, Commun. Theor. Phys. **45**, 151 (2006).

[67] Z. H. Chen, Z. B. Yang, C. G. Liao, and C. L. Luo, Opt. Commun. **284**, 297 (2011).

[68] S. B. Zheng, Chin. J. Laser B **7**, No. 6, 534 (1998).